

Reference #: 923239
Report Date: 24 Sep 2015

Date Received: 16 Sep 2015

Referring Veterinarian; DR. GLENN FAHNESTOCK EASTVIEW VETERINARY CLINIC P.O. BOX 237

PENN YAN, NY 14527 UNITED STATES Patient ID:

15091601643

Radiography Date:

16 Sep 2015

Owner/Responsible Person:

KAY FRANK

Patient: Patient Name: CALL Species: CANINE Reg. Name: Breed: **GERMAN SHEPHERD** Reg. #: Tatloo: Date of Birth: 6 Nov 2014 Age: 10 mo. Weight: 73 lbs. Microchip: Gender:

			RESULTS				
	Distraction Index (DI) 0.38		DI is greater than 0.30 with no radiographic evidence of OA. There is an				
LEFT	Osteoarthritis (OA)	None	increasing risk of developing OA as the DI Increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.				
ï	Cavilation	No					
	Other Findings	Not Applicable					
	Distraction Index (DI)	0.36	Dt is greater than 0.30 with no radiographic evidence of OA. There is an				
RIGHT	Osteoarthritis (OA) None		increasing risk of developing OA as the DI increases; low risk when DI is close to 0.30, high risk when DI is close to 0.70 or above.				
	Cavitation	No					
	Other Findings	Not Applicable					

Please note that the PennHIP DI is a measure of hip joint laxity, it does not allude to a "passing" or "failing" hip score.

LAXITY PROFILE RANKING

The laxity profile ranking is based on the hip with the greater laxity (DI). This interpretation is based on a cross-section of 12,012 CANINE animals of the GERMAN SHEPHERD breed. The median DI for this group is 0.40.

					Percentiles					
	90th	80th	70th	នក្សា	50th	40th	30th	20th	10th	
> 90th					Median					< 10th

The chart above indicates the ranking of your animal's passive hip laxity (DI) in relation to all CANINE animals of the GERMAN SHEPHERD breed in our database. This result means that 1) your animal's hips are tighter than approximately 60% of this group of animals (alternatively, 40% of the group has tighter hips than your animal), and 2) your animal's hip laxity is in the tighter half of the laxity profile. Breed-specific evaluations are analyzed semi-annually. Consequently, the average laxity and range of laxity for any given group will change over time.

PennHIP does not make specific breeding recommendations. Selection of sire and dam for mating is the decision of the breeder.

NOTE: As a minimum breeding critarion, we propose that breeding stock be selected from the population of animals having hip laxity in the tighter half of the breed (to the left of the median mark on the graph). Higher selection pressure equates to more rapid expected genetic change per generation.

By implementing selection based on passive hip laxity, we expect the breed average DI over the years to move toward tighter hip configuration, meaning lower hip dysplasia susceptibility. The PennHIP database permits scientific adjustment of criteria to reflect these shifts; the average laxity and range of laxity for a particular breed will change over time.

ANTECH Imaging Services / 17672-B Cowen Avenue / Irvine, CA 92614 677-727-6800 / Fax: 877-870-4890 www.antechimagingservices.com/pennhip